



FACT SHEET

Amebiasis

What is it?

Amebiasis is an intestinal infection caused by a parasite, *Entamoeba histolytica*. The parasite produces cysts (eggs) which are passed from the body in the stool (bowel movements).

Symptoms

- Fever
- Chills
- Diarrhea (sometimes bloody or with mucus).
- Some people may have only mild abdominal discomfort or no symptoms at all.
- The symptoms start 2-4 weeks or even months after infection with a cyst.

Potential Complications

- In rare cases the parasite may invade the liver, lung, brain or perforate the colon causing death.

How is it spread?

- The parasite is spread from one person to another directly or via contaminated objects or food or water if an infected person does not wash his or her hands well after going to the bathroom. When others touch the objects or eat contaminated food they get the cysts on their hands and into their mouths. When they swallow the cysts they get the infection.
- People are infected as long as the cysts are shed in the stool

Diagnosis and Treatment

- Diagnosis is done by examining a stool sample under a microscope for the presence of the parasite or its cysts.
- People who have Amebiasis are usually treated with prescription medication. There are no over-the-counter medications that will cure this infection.

Prevention

- Practice good hand washing habits after going to the bathroom, before preparing meals and before eating.
- Practice good hand washing after changing diapers.

Additional Information

- People who travel to areas of the world which do not have sanitary water supply are at greater risk for getting amebiasis.
- Men who have sex with men are also a greater risk of getting amebiasis through sexual contact.

Report all cases to El Paso Department of Public Health by calling 915-212-6520