



FACT SHEET

West Nile Virus

What is it?

- West Nile virus (WNV) cause West Nile fever and West Nile meningitis, encephalitis, and West Nile poliomyelitis. It can infect people, birds, mosquitoes, horses and other animals.

What are the symptoms?

- Most people who are infected with the West Nile virus will not get sick. In those who do become ill, symptoms usually begin between 3 and 14 days after being bitten by an infected mosquito.
- After 1 in 5 infected persons will develop West Nile fever, a flu-like illness lasting a few days to several weeks.
- In addition to fever symptoms include:
 - Fatigue
 - Headache
 - Loss of appetite
 - Muscle aches
 - Nausea
 - Rash
 - Vomiting
 - Swollen glands
- About 1 in 150 infected people will have one of the more severe and potentially fatal forms of disease (West Nile meningitis, encephalitis, or West Nile poliomyelitis) that can include symptoms of West Nile fever and:
 - Severe headache
 - Neck stiffness
 - Sensitivity to light
 - Decreased level of alertness
 - Muscle weakness, tremors, or paralysis
 - Convulsions or coma

How is it spread?

- West Nile virus is spread to people mainly by the bite of an infected mosquito. Mosquitoes become infected after feeding on birds that carry the virus. WNV is not spread by direct contact with infected animals or people, but care should always be used when handling sick or dead animals. WNV may be spread via blood transfusion and organ transplantation. Blood and organ donations are routinely tested and infected donations are not used.

Who is at risk?

- The risk of getting West Nile virus is very low, but anyone who is bitten by infected mosquitoes can

be infected. Older people over 50 years of age have a higher risk of having more serious symptoms.

Diagnosis and treatment

- West Nile virus in humans is diagnosed by testing blood and/or CSF (cerebrospinal fluid from a spinal tap)
- Contact your healthcare professional if you have symptoms of a possible WNV infection, especially if you recently had mosquito bites.
- There is no vaccine for humans or specific treatment other than supportive care.

Prevention

- Empty anything outdoors that holds standing water – any standing water can serve as a mosquito breeding site.
- Change water in birdbaths, fountains, wading pools and animal troughs weekly.
- Make sure that roof gutters drain properly.
- Fix leaky outdoor faucets and sprinklers.
- Make sure windows and door screens are “bug-tight” – if not, repair or replace them.
- Stay indoors at dawn and dusk when mosquitoes are the most active.
- Wear protective clothing (long sleeves, long pants and a hat) when going into mosquito-infested areas.
- Use mosquito repellent when necessary, and carefully follow the directions on the label. Effective repellents contain DEET, picaridin, or oil of lemon eucalyptus.
- Assist elderly neighbors and relatives in these tasks to help keep them safe from mosquito bites.

What should I do if I find a dead bird?

- West Nile virus infects many wild birds including crows, jays and ravens. Increasing number of dead birds can be a sign that WNV is present in the community. As with any dead animal, avoid any direct contact. Wear gloves or use a shove to move the bird. Dead birds can be disposed of by putting them in a plastic bag and then into the garbage.

Report all cases to El Paso Department of Public Health by calling 915-212-6520

